

Time varying gauge couplings and di-nucleon states

Malcolm Fairbairn

*Service de Physique Théorique, CP225,
Université Libre de Bruxelles, B-1050 Brussels, Belgium*

In this talk we discuss the effect upon the di-proton, di-neutron and deuteron of a time variation of the QCD gauge coupling. We describe how a time evolution of the size of extra dimensions can give rise to such a variation and show how this is related to the recent results on possible time variation of the electromagnetic fine structure constant. Based on work in collaboration with Tom Dent.

1 Introduction

Webb et al have observed the absorption of light from quasars by intervening intergalactic clouds of matter at high redshifts¹. By studying the relative position of absorption lines in these clouds, they have found that there appears to be a variation in the electromagnetic fine structure constant α_{em} at high redshift.

Such a variation is naturally accommodated in higher dimensional theories such as string or M-theory since in these models the value of the gauge couplings are set by the size of the compactified directions. Cosmological evolution of the higher dimensions would then lead to time variation of the gauge coupling of the grand unified theory. One would expect this to lead to a variation of all of the gauge couplings in the standard model including the QCD coupling.

The renormalisation group running of the QCD coupling constant α_3 means that the accompanying fractional variation in the characteristic energy scale of hadronic physics Λ_{QCD} would be much larger than that for α_{em} . It is natural therefore to investigate as to whether or not the observed variation is consistent with the assumptions of gauge coupling unification and physics in the hadronic sector.

After briefly discussing the results, we talk more about the relationship between the size of higher dimensions and the gauge couplings in the 4D theory in one particular model. Next, we describe how variations in α_{em} lead to variations in Λ_{QCD} and the implications of this for the binding of the two nucleon states, the deuteron, di-neutron and di-proton. We then show that this leads to constraints on the variation of α_{em} from the variation of the binding energy of the deuteron during nucleosynthesis.

2 The observations

As we have said, Webb et al¹ have measured α_{em} at high redshifts by observing the absorption lines in the light from high redshift quasars. This absorption occurs in clouds of intervening intergalactic dust. When trying to understand the origin of a time variation in α_{em} , it is useful to know a tiny part of the atomic physics that allow one to probe such a variation.

Upon solution of Schrodinger's equation in atomic physics, the n th electron energy level of a given atom is proportional to

$$E \propto -\frac{me^4}{2\hbar^2} \frac{Z^2}{n^2} \quad (1)$$

where m is the electron mass, e is it's charge and Z is the number of protons in the nucleus. The relativistic correction to this energy level has the following proportionality²

$$\Delta \propto \frac{me^4}{2\hbar^2} \frac{Z^2}{n^2} \frac{(Z\alpha_{em})^2}{n}. \quad (2)$$

Comparison of the previous two equations is the key to understanding why observation of spectral lines in high redshift objects is an extremely sensitive probe of the variation of the gauge couplings. Typically, if one was trying to determine the energy of a particular spectral line in a high redshift object, there would be many systematic errors due to uncertainties in one's knowledge of the redshift or proper motion of the object. However, these systematic errors fall out of the analysis if one deduces α_{em} by comparing a number of lines within the same cloud. It is also interesting to note that such measurements are not sensitive to a time variation in the mass of the electron, since this would simply act like an apparent change in the redshift of the system.

It is then possible to compare the observed lines to those predicted numerically to see if the best fit is with α_{em} as we know it today, or with a different value of α_{em} . The analysis of Webb et al¹ shows that for redshifts of approximately $z \sim 1 - 3$ the observed spectral lines are statistically significantly better fit by the numerical models if one assumes that α_{em} was smaller by about 1 part in 10^5 at those times.

It is therefore interesting to consider the effect of this variation assuming that the gauge fields are part of some larger GUT symmetry and that variation of one gauge coupling leads to variation of all three gauge couplings.

3 GUT theories and higher dimensions

The observed unification of the gauge couplings at 10^{16} GeV in the supersymmetric (SUSY) version of the standard model strongly³ suggests that both QCD and the electroweak theory are parts of a larger gauge symmetry broken at high energy. In this section we show how the coupling of the GUT gauge field is related to the size of extra dimensions in one particular model, namely Heterotic M-theory. We then report how this all relates to α_{em} and the hadronic sector following the analysis of our paper⁴.

3.1 Some (M)otivation

We know that in order to have general relativity and some unified gauge theory with field strength F containing the standard model, the Lagrangian of the effective four dimensional theory obtained by compactification of some particular string/M-theory must look like

$$L_{eff} = - \int d^4x \sqrt{g} \frac{1}{16\pi G} R - \int d^4x \sqrt{g} \frac{1}{16\pi\alpha_{GUT}} F^2. \quad (3)$$

As we mentioned earlier, the values of the gauge and gravitational couplings α_{GUT} and G are set by the size of the extra dimensions. This arises because the volume of the higher dimensional space in which each particular field propagates re-scales the coupling of that field upon compactification.

For example in Heterotic M-theory, the GUT gauge field comes from the E_8 gauge field in the 11 D theory⁵. This gauge field exists only in the uncompactified directions and in the 6

extra directions which are compactified upon a Calabi-Yau manifold. The coupling in 4D is given by

$$\alpha_U = \frac{2\pi^{2/3}\kappa^{4/3}}{V} \quad (4)$$

where κ^2 is the 11D Newton's constant and V is the volume of the Calabi-Yau. As usual, gravity propagates in all space-time directions including the 11th dimension and the effective Newton's constant in 4D is given by

$$G = \frac{\kappa^2}{16\pi^2 V \rho} \quad (5)$$

where ρ is the size of the 11th (orbifolded) dimension. So it is possible to have variations in the 4D strength of gravity if ρ evolves over time and in the strength of both gravitational and gauge interactions if V evolves over time.

There have been various suggestions as to the mechanisms responsible for stabilising V and ρ at least in weakly coupled heterotic string theory. Perhaps the most well known are non-perturbative corrections to the scalar potential for these fields due to the condensation of the fermionic superpartners of gauge bosons^{6,7}, but the problem is not completely solved and becomes more complicated in the strong coupling regime.

3.2 Other aspects of time varying extra dimensions

In all compactifications of higher dimensional theories, if the sizes of the extra dimensions are allowed to possess dynamics they give rise to scalar fields in the 4D theory with cosmological equations of motion of the usual form

$$\ddot{\phi} + 3H\dot{\phi} + m^2\phi = 0 \quad (6)$$

where m is the second derivative of the potential in which the field is rolling and $H = \dot{a}/a$ is the Hubble expansion factor. Variation of the size of the higher dimensions over time leads to variations in these scalars ϕ and to the variation of our effective 4D gauge couplings. Now the observed variation of the fine structure constant is of order $\Delta\alpha_{em}/\alpha_{em} \sim 10^{-5}$ at a redshift of order $z \sim 1$ ¹. Nucleosynthesis occurs at redshifts of order $z \sim 10^8 - 10^{10}$ and the variation in gauge couplings at that time is certainly constrained to at least $\Delta\alpha_{em}/\alpha_{em} < 1$. If this were not the case, it would not be possible to obtain a primordial ratio of Helium and Hydrogen that matches that observed in regions of the universe where no nuclear processing has occurred since nucleosynthesis^{8,9}. Therefore if one is to take seriously dynamics of the form of equation (6) it is either necessary to cook up a very original potential, or to assume that the slope of the potential is extremely small so that equation $\dot{\phi}$ is damped and the ϕ is frozen at early times when

$$m^2 \ll H^2. \quad (7)$$

The variation of the gauge coupling would then begin later on when the expansion drops to the point where $H^2 \sim m^2$ and the field ϕ is no longer frozen. This leads to a fine tuning problem in explaining why m is so small. Also one might expect a very small mass for such a field to lead to dangerous light modes¹⁰ and/or have associated with it a particle leading to new long range forces¹¹. Therefore a variation at such a late time, if proved to be correct, would create so many problems for our existing model that it would give us a great deal of information about not just unification but many other aspects of particle physics beyond the standard model.

3.3 $\Lambda_{QCD} = f(\alpha_{em})$ in GUT theories

The relationship between Λ_{QCD} and the gauge coupling in the grand unified theory is obtained by considering the relationship between Λ_{QCD} and the QCD gauge coupling α_3 at energies just

above the masses of the light quarks. Then one can find the relation between α_3 at these low energies and at the energy scale of unification¹² M_U to obtain

$$\Lambda_{QCD} = M_{SUSY}^{\frac{2}{3}} M_U^{\frac{1}{3}} \left(\frac{m_c m_b m_t}{M_{SUSY}} \right)^{\frac{2}{27}} \exp \left(\frac{-6\pi}{27\alpha_3(M_U)} \right) \quad (8)$$

where m_i is the mass of the i quark and we have assumed that all the SUSY particles have the same mass which is equal to the mass of SUSY breaking M_{SUSY} . Next, by running the electroweak gauge couplings α_1 and α_2 down from M_U to low energies we obtain the relation

$$\left. \frac{\Delta\Lambda_{QCD}}{\Lambda_{QCD}} \right|_{M < m_e} \sim 34 \left. \frac{\Delta\alpha_{em}}{\alpha_{em}} \right|_{E \ll m_e} \quad (9)$$

the linearity being applicable for variations in α_{em} less than a few parts in 10^3 .

It is worth pointing out that at high energies, and therefore high temperatures in the early universe, one would expect the same renormalisation to change the value of the gauge couplings and hence α_{em} . How do we know this is not responsible for the possible time variation of α_{em} ? Well apart from the fact that the spectral lines correspond to a jump in electron energy and therefore are more like a tunnelling process than something like compton scattering, the variation goes the wrong way and one would expect no renormalisation of α_{em} below the electron mass (1 MeV, $z \sim 10^{10}$) anyway.

4 The effect of varying gauge couplings on di-nucleon states

4.1 The internuclear pion potential

If one ignores the small contribution due to the bare quark masses and the interquark electromagnetic binding energy, the nucleon mass M_N and the masses of mesons such as the sigma and omega meson scale with Λ_{QCD}

$$M_N \propto m_\omega \propto m_\sigma \propto \Lambda_{QCD}. \quad (10)$$

This is not true for the pion mass which arises as the Goldstone boson of spontaneously broken chiral symmetry. Goldstone bosons of exact symmetries are massless, but the fact that the underlying chiral symmetry of the QCD Lagrangian is already slightly broken by the presence of small but non zero quark masses leads to a pion mass which scales as (see e.g. ¹³)

$$m_\pi \propto (m_q \Lambda_{QCD})^{\frac{1}{2}} \quad (11)$$

where m_q is the average mass of the light quarks. Here we have also assumed that the energy scales of confinement and chiral symmetry breaking coincide as suggested by simulations on the lattice⁴. The exchange of pions contributes to the internuclear potential as

$$V(r) = -\frac{f^2}{4\pi} \frac{e^{-m_\pi r}}{r} \quad (12)$$

where the strength of the force is given by

$$f^2 = \left(\frac{g_\pi^2 m_\pi^2}{4M_N^2} \right) \propto \frac{m_q}{\Lambda_{QCD}} \quad (13)$$

where g_π is the pion nucleon coupling and M_N is the nucleon mass⁴.

4.2 The stability of the di-nucleons

The fact that the di-proton and the di-neutron are not bound is essential for the universe to evolve the way it has done. This is because the existence of a stable di-proton would lead to a rapid channel for the burning of hydrogen in stars. If the di-proton was stable, it would not be possible for stars to slowly burn hydrogen on the main sequence for long periods of time so the fact that the di-proton is unbound is essential for our understanding of stellar evolution. Since the internuclear potentials in the di-proton and di-neutron systems are described by the same nuclear potential (12), the only difference between the two binding criteria arises from electromagnetic effects which are very much smaller than the precision of the results in this paper.

We can use the potential (12) to calculate the amount that one would have to change the ratio between m_q and Λ_{QCD} in order to stabilise the di-neutron or di-proton system. We use a trial wavefunction of the form

$$\psi(r) = e^{-1/m_\pi r} e^{-bm_\pi r} \quad (14)$$

so that at small r the second derivative of the wavefunction depends mostly upon the potential whereas at large r it depends mostly upon the energy, denoted by the variational term b . Then we can write




Figure 1: Stability of di-nucleon system. Dark region is bound, light region unbound.

$$v = \frac{f^2}{4\pi} \frac{2M_N}{m_\pi} \propto \frac{m_q}{\Lambda} \quad (15)$$

Once we have evaluated v , we can see from figure 1 that in order for the di-nucleon system to remain unbound, $\Delta v/v < 2.6$. This corresponds to a very large change in Λ_{QCD} and the corresponding change in α_{em} is about -5%. A similar analysis for the deuteron leads to a criterion for un-binding of about a +2% increase in α_{em} .

4.3 Nucleosynthesis Constraints

We have placed constraints on the amount by which it is possible to vary α_{em} so that the corresponding variation in Λ_{QCD} is small enough to be consistent with the stability of the deuteron and the instability of the di-nucleon. Of course, without saturating these bounds, a variation in the ratio between the quark masses and Λ_{QCD} will lead to a change in the binding energy of the deuteron. This will have implications for the production of Helium during nucleosynthesis.

When the universe is hotter than about 1 MeV, neutrons and protons are continuously interconverting via the weak interactions

$$\begin{aligned} n + e^+ &\leftrightarrow p + \bar{\nu}_e \\ n + \nu_e &\leftrightarrow p + e^- \\ n &\leftrightarrow p + e^- + \bar{\nu}_e. \end{aligned}$$

with the rate of these reactions being given approximately by

$$\Gamma_{pe \rightarrow n\nu} \approx G_F^2 T^5 \quad (16)$$

Now when the Hubble expansion $H = 1.66\sqrt{g_*}GT^2$ becomes larger than this rate the reactions freeze out. This occurs at a temperature

$$T_o \approx \frac{G^{1/6}}{G_F^{2/3}} \quad (17)$$

and the ratio of neutron to proton number density (n/p) which one finds at this temperature is frozen, only changing via the beta decay of the neutron.

$$(n_o/p_o) = e^{-(M_n - M_p)/T_o} \quad (18)$$

Later at $T_{He} \sim B_{deut}$, the deuteron can survive without becoming photo-dissociated by the hot plasma and the following Helium forming reactions are able to take place

$$\begin{aligned} &H^2(H^2, n)He^3(H^2, p)He^4 \\ &H^2(H^2, p)H^3(H^2, n)He^4 \\ &H^2(H^2, \gamma)He^4 \end{aligned} .$$

Essentially all (99.99%) of the neutrons around at this temperature form helium via deuterium. The mass fraction of Helium is then given by

$$Y = X_4 = \frac{2(n/p)}{1 + (n/p)} \Big|_{T_{He}} ; \quad X_A = \frac{n_A A}{n_N} \quad (19)$$

Now as the neutrons decay, the neutron proton ratio changes

$$(n/p) = \frac{\exp(-t/\tau)}{(p_o/n_o) + 1 - \exp(-t/\tau)} \quad (20)$$

where τ is the neutron lifetime, so the time between weak freeze out and the temperature at which Helium can form is essential in determining the resulting mass fraction of Helium in the universe. Since this temperature is determined by the binding energy of the deuteron, the process is extremely sensitive to variations in the internuclear potential and consequently the

gauge couplings. One finds that in order to agree with the observed primordial mass fraction of Helium the binding energy of the deuteron is constrained to lie within

$$-0.20 \leq \frac{\Delta B_{deut}}{B_{deut}} \leq +0.13 \quad (21)$$

which corresponds to a variation in Λ_{QCD} of

$$+0.04 \geq \frac{\Delta \Lambda_{QCD}}{\Lambda_{QCD}} \geq -0.04. \quad (22)$$

If we calculate what variation in the gauge couplings can give rise to this variation in Λ_{QCD} we see that

$$-0.005 \leq \left. \frac{\Delta \alpha_3}{\alpha_3} \right|_{1 \text{ TeV, no SUSY}} \leq 0.005 \quad (23)$$

for the variation in α_3 at 1 TeV, assuming that is where the gauge couplings are set. If we consider the more conventional case with SUSY and gauge coupling unification at 10^{16} GeV, we can find out exactly what this means in terms of the variation of α_{em}

$$\left| \frac{\Delta \alpha_{em}}{\alpha_{em}} \right| \leq 10^{-3}. \quad (24)$$

This shows us that the constraint on α_{em} is very strong around nucleosynthesis, so if it is varying today, it cannot have varied much more rapidly in the early universe.

5 Conclusions

We have shown that a variation of α_{em} implies a much larger variation in Λ_{QCD} if the gauge couplings are unified at very high energies. This leads to constraints upon the variation in α_{em} due to the fact that we must maintain the stability of the deuteron and the instability of di-proton at all times since nucleosynthesis. These constraints are typically of the order for a few percent. We have also shown that it is possible to place new constraints upon the variation of α_{em} at nucleosynthesis since quite a small variation in α_{em} would give rise to a large variation in the primordial Helium abundance if one assumes gauge coupling unification.

Acknowledgments

This is based on work in collaboration with Thomas Dent, and I am grateful for conversations with Jean-Marie Frère.

References

1. J. K. Webb, M. T. Murphy, V. V. Flambaum, V. A. Dzuba, J. D. Barrow, C. W. Churchill, J. X. Prochaska, and A. M. Wolfe, *Phys. Rev. Lett.* **87**, 091301 (2001)
2. V. A. Dzuba, V. V. Flambaum, and J. K. Webb, *Phys. Rev. Lett.* **82**, 888 (1999)
3. D. M. Ghilencea and G. G. Ross, *Nucl. Phys. B* **606**, 101 (2001)
4. T. Dent and M. Fairbairn, [arXiv:hep-ph/0112279]
5. E. Witten, *Nucl. Phys. B* **471**, 135 (1996)
6. A. Font, L. E. Ibanez, D. Lust and F. Quevedo, *Phys. Lett. B* **245**, 401 (1990)
7. T. Barreiro, B. de Carlos and E. J. Copeland, *Phys. Rev. D* **57**, 7354 (1998)
8. B. A. Campbell and K. A. Olive, *Phys. Lett. B* **345**, 429 (1995)
9. B. E. J. Pagel, *Phys.Rep.* **333**, 433, (2001)

10. B. de Carlos, J. A. Casas, F. Quevedo and E. Roulet, *Phys. Lett. B* **318**, 447 (1993)
11. G. R. Dvali and M. Zaldarriaga, *Phys. Rev. Lett.* **88**, 091303 (2002)
12. Bailin, D. and Love, A., Introduction to Gauge Field Theory, IOP publishing, (1993),
Supersymmetric Gauge Field Theory and String Theory, IOP publishing, (1996)
13. S. Weinberg, "The quantum theory of fields. Vol 2: Modern applications", Cambridge University Press (1996)